


## How does GENGIGEL® work?

### THE 3 SYNERGISTIC EFFECTS OF GENGIGEL®

- ☛ Regulates periodontal inflammatory processes.<sup>3</sup>
- ☛ Reduces gum inflammation and bleeding.<sup>3</sup>

- ☛ Accelerates repair and cicatrisation mechanisms.<sup>4</sup>



- ☛ Act as barrier to macromolecules and foreign bodies.<sup>4</sup>

☛ **GENGIGEL®** by the virtue of its high viscosity and elasticity -

- Modulates inflammation.<sup>5</sup>
- Possesses bacteriostatic effect and anticandidal activities.<sup>4,6</sup>
- Antiedematous effect.<sup>5</sup>
- Improves tissue regeneration.<sup>5</sup>
- Stimulates angiogenesis.<sup>4,5,7</sup>
- Accelerates bone regeneration.<sup>8</sup>

## GENGIGEL® SKUs

### General Range



#### GENGIGEL® Gel 1ml Sachets (0.2% HA) Pack of 100 sachets

##### Direction:

Apply a film of gel onto the gingival tissue, massaging gently with clean fingers to aid its correct distribution on the mucosal surface.



#### GENGIGEL® Hydrogel Mouthrinse 150ml (0.025% HA)

##### Direction:

Rinse with 10ml (one dose) for 1-2 minutes, 3 times a day. The compound is ready to use, do not dilute.



#### GENGIGEL® Gel 20ml Tube (0.2% HA)

##### Direction:

Apply a film of gel onto the gingival tissue, massaging gently with clean fingers to aid its correct distribution on the mucosal surface.



#### GENGIGEL® Gel 20ml Spray (0.01% HA)

##### Direction:

1 to 2 puffs on inflammatory or traumatised tissue. Each puff guarantees homogeneous constant distribution of the liquid on the mouth surface.

### Professional Range



#### GENGIGEL® PROF FLUID 500ml (0.1% HA)

##### Indications:

1. Provides pain relief.
2. Promotes and accelerates tissue reconstruction after any traumatic condition (surgical intervention, implants, tooth extraction, root scaling, etc).
3. Promotes and accelerates the anti-oedematous and anti-inflammatory processes.
4. Indicated for gingival pockets, all forms of periodontitis, including serious forms, at the perioperative stages and in any situation in which wounds penetrate the dermis.



#### GENGIGEL® PROF SYRINGES

##### Pack of 4 X 1ml Syringes (0.8% HA)

##### Indications:

1. Promotes and accelerates tissue reconstruction after any traumatic condition (surgical intervention, implants, tooth extraction, root scaling, etc).
2. After non-surgical thorough debridement reduces pocket depth.
3. Promotes and accelerates the anti-oedematous and anti-inflammatory processes.
4. Reduces bleeding.
5. Indicated for all forms of periodontitis, including serious forms, at the perioperative stages and in any situation in which wounds penetrate the dermis.

Store at 5°C -37°C

Do not consume any food or drinks for 30 minutes after **GENGIGEL®** application





## GENGIGEL® A FOUNDATION FOR DENTAL OUTCOME

- Reduces inflammation and bleeding of gums.<sup>2</sup>
- Accelerates tissue proliferation and healing of gum injury.<sup>2</sup>
- Maintains healthy gingival tissue.<sup>2</sup>
- Indicated across Gingivitis to Periodontal management continuum.<sup>12</sup>
- Contains xylitol and does not cause caries. It is suitable for diabetics.<sup>12</sup>
- Obtained by biotechnological synthesis.<sup>12</sup>
- Biomimetic Hyaluronic Acid.<sup>12</sup>
- Medical device Class B certified and approved by EU and FDA.<sup>12,13,14</sup>
- Safe to use in young children, pregnant women and elderly patients.<sup>12</sup>

### References

**1.** Periodontal status and provision of periodontal services in Malaysia Trends and way forward, Mohd-Dom TN, et al. MJPHM, 2013; 13(2): 38-47.; **2.** Hyaluronic Acid: A Boon in Periodontal Therapy. Parveen D, et al. N Am J Med Sci. 2013;5(5):309-315. **3.** The clinical application of hyaluronic acid in gingivitis therapy. Pistorius A, Martin M, Willershausen B et al. Quintessence Int 2005;36:531-538. **4.** A promising mediator for periodontal regeneration. Bansal J, Kedige SD, Anand S. Hyaluronic acid: Indian J Dent Res 2010;21:575-578. **5.** A model for the role of hyaluronic acid and fibrin in the early events during the inflammatory response and wound healing. Weigel PH, Fuller GM, Le-Boeuf RD. J Theor Biol 1986;119(2):219-234. **6.** Potential role of high molecular weight hyaluronan in the anti-Candida activity of human oral epithelial cells. Sakai A, Akifusa S, Itano N et al; Med Mycol Feb 2007;45(1):73-79. **7.** Hyaluronic acid and periodontitis. Sukumar S, Drízhali I; Acta Medica (Hradec Kralove) 2007;50:225-258. **8.** Stimulation of Osteoinduction in Bone Wound Healing by High-Molecular Hyaluronic Acid Sakasi T, Watanabe C., 1995;16:9-15. **9.** Gengigel Product Monograph. **10.** Application of Gengigel in the Treatment of Gingival Inflammation M. Popovska, R. Kievski, B. Angelovski, Z.Belazelkoska, E. Stefanovska, L. Kanurkova; Balk J Stom, 2012; 16:43-45. **11.** Hyaluronic Acid as an Adjunct After Scaling and Root Planing , Sigrun Eick, Antonio Renatus, DMD, Melanie Heinicke, Wolfgang Pfister, Professor, Stefan-Ioan Stratul, Professor, Holger Jentsch, Journal of Periodontology; 2012 DOI: 10.1902/jop.2012.120269. **12.** Gengigel Product Insert **13.** Gengigel Medical Device (MDA) Registration Certification. **14.** Gengigel Food Drug Administration Certification.

Please consult your healthcare professional for further information.

**pharmaniaga**<sup>®</sup>  
passion for patients

**PHARMANIAGA MARKETING SDN BHD (118254-D)**  
No. 7, Lorong Keluli 1B, Kawasan Perindustrian Bukit Raja Selatan, Seksyen 7,  
40000 Shah Alam, Selangor Darul Ehsan, Malaysia.  
Tel : 603-3342 9999 Fax : 603-3344 2222 Customer Care Line : 1-800-888-313  
Email : customercare@pharmaniaga.com Website : www.pharmaniaga.com

 **RICERFARMA** S.R.L.  
[www.ricerfarma.com](http://www.ricerfarma.com)

Via Egadi, 7 - 20144 Milano - Italy  
Email : info@ricerfarma.com

**For Healthcare professional use only**